

Example 1

Let $\{x_n\}_{n=1}^{\infty}$ be the sequence $1, -2, 3, -4, 5, -6, 7, -8, 9, -10, \dots$ so it is given by $x_n = (-1)^{n+1}n$.

Example 2

Let $\{x_n\}_{n=1}^{\infty}$ be the sequence $1, 2, 3, 1, 5, 6, 1, 8, 9, 1, 11, 12, 1, 14, 15, 1, 17, 18, 1, \dots$

- Roughly speaking, a **subsequence** of a sequence $\{x_n\}_{n=1}^{\infty}$ is obtained by choosing infinitely many values from the given sequence, where each successive choice is made by taking a larger index (i.e. looking farther out in the sequence).

Exercise.

- Give a few reasons you know that the sequences in Example 1 and Example 2 are divergent.
- Write down a few examples of subsequences of the sequence in Example 1.
- Write down a few examples of subsequence of the sequence in Example 2.
- Do either of the sequences in Example 1 or Example 2 have the property that it has a convergent subsequence?

Example 3.

Let $\{x_n\}_{n=1}^{\infty}$ be the sequence $x_n = \cos n$.

Exercise.

- a) Write out the first several terms of the sequence in Example 3.
- b) Can you tell whether or not the sequence converges?
- c) Can you tell whether or not the sequence has a convergent subsequence?

Example 4.

Let's define a sequence $\{x_n\}_{n=1}^{\infty}$ as follows: For each $n \in \mathbb{N}$, let x_n be any number at all between 0 and 100.

Exercise.

a) Is the sequence in Example 4 convergent?

b) If we try to answer the question

“does the sequence in Example 4 have a convergent subsequence?”,

why does this question appear to be harder to answer than the same question with Example 1 or with Example 2?

- In this section our interest is in deciding for a given sequence whether or not we can be certain that the sequence has a convergent subsequence.
- We won't be able to decide completely, but

the **Bolzano – Weierstrass theorem** gives a sufficient condition on a given sequence which will guarantee that it has a convergent subsequence.

- So the theorem will guarantee that if the given sequence satisfies the hypothesis of the Bolzano-Weierstrass theorem, then we know for certain that the sequence has a convergent subsequence, even if we don't know how to explicitly write that subsequence down.

Before we state the theorem, let's first give a formal definition of subsequence of a sequence.

Definition

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence. Let $n_1 < n_2 < n_3 < n_4 < \dots$ be a strictly increasing sequence of natural numbers. Let $\{y_k\}_{k=1}^{\infty}$ be the sequence defined by $y_k := x_{n_k}$. Then the sequence y_k is called a **subsequence** of the sequence x_n .

Alternate Definition

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence. We say that sequence $\{y_n\}_{n=1}^{\infty}$ is a **subsequence** of $\{x_n\}_{n=1}^{\infty}$ if there exists a function $\phi : \mathbb{N} \rightarrow \mathbb{N}$ which is strictly increasing such that $y_n = x_{\phi(n)}$ for every n .

Exercise.

Consider the sequence of Example 2: 1, 2, 3, 1, 5, 6, 1, 8, 9, 1, 11, 12, 1, 14, 15, 1, 17, 18, 1, \dots

- Using the alternate definition, what subsequence do we get if we take $\phi(n) = 2n$?
- What choice of $\phi(n)$ gives the subsequence 1, 1, 1, 1, \dots ?

Alternate Definition

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence. We say that sequence $\{y_n\}_{n=1}^{\infty}$ is a **subsequence** of $\{x_n\}_{n=1}^{\infty}$ if there exists a function $\phi : \mathbb{N} \rightarrow \mathbb{N}$ which is strictly increasing such that $y_n = x_{\phi(n)}$ for every n .

Exercise.

Suppose $\{x_n\}_{n=1}^{\infty}$, $\{y_n\}_{n=1}^{\infty}$, $\{z_n\}_{n=1}^{\infty}$ are three sequences for which $\{y_n\}_{n=1}^{\infty}$ is a subsequence of $\{x_n\}_{n=1}^{\infty}$ and $\{z_n\}_{n=1}^{\infty}$ is a subsequence of $\{y_n\}_{n=1}^{\infty}$. Prove that $\{z_n\}_{n=1}^{\infty}$ is a subsequence of $\{x_n\}_{n=1}^{\infty}$.

Is it possible that boundedness or unboundedness has something to do with whether or not a sequence has a convergent subsequence?

Exercise.

Write down two sequences with the following properties:

- Both sequences consist entirely of positive numbers.
- Both sequences are unbounded.
- The first sequence does not have a convergent subsequence, but the second sequence does have a convergent subsequence.

Exercise.

So what does the above exercise tell you about what the Bolzano-Weierstrass theorem can definitely not be?

Exercise.

- a) Recall that Example 4 had a sequence $\{x_n\}_{n=1}^{\infty}$ consisting of randomly chosen numbers between 0 and 100. Why is it plausible that this sequence has a convergent subsequence?
- b) This example suggests a conjecture as to what might be a sufficient condition to guarantee that a sequence must have a convergent subsequence. What is that condition?

Theorem (Bolzano-Weierstrass)

Let $\{x_n\}_{n=1}^{\infty}$ be any bounded sequence. Then $\{x_n\}_{n=1}^{\infty}$ has a convergent subsequence.

Comments on the proof

- It is sufficient to show that the sequence has a Cauchy subsequence. The result will then follow from the completeness axiom of \mathbb{R} .
- This is done using the method of interval halving. We show how to construct a nested decreasing sequence of intervals $I_n = [a_n, b_n]$ such that $(b_n - a_n) \rightarrow 0$ and a subsequence y_n of the original sequence such that $y_n \in I_n$ for each n . Then y_n is automatically Cauchy.
- Let $I_1 = [a_1, b_1]$ be any closed interval which contains the entire sequence. This is possible since the sequence is bounded. Let $n_1 := 1$ and $y_1 := x_{n_1} = x_1$.
- The key idea which makes the proof work is the fact that
either the left half or the right half of I_1 must contain infinitely many terms of the sequence,
so we let I_2 be the half which does contain infinitely terms of the sequence, and we let y_2 be a specific term x_{n_2} of the sequence which is in I_2 .
- After that, we apply the same idea on I_2 to produce an interval I_3 and a number y_3 . The rest of the intervals and points are obtained in a similar manner.

Theorem (Bolzano-Weierstrass)

Let x_n be any bounded sequence. Then x_n has a convergent subsequence.

Exercise.

Use the comments on the previous slide to write the proof of the Bolzano-Weierstrass Theorem

Example

Revisit Examples 3 and 4. Explain how we know that those sequences must have convergent subsequences.

Exercise.

Formulate a theorem like the Bolzano-Weierstrass Theorem which applies to all sequences, including sequences which are unbounded. Thus it should read

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence such that *(you fill in the blank)*.

Then $\{x_n\}_{n=1}^{\infty}$ has a convergent subsequence.