

## 1.6. One-sided Limits and Continuity

### One-sided Limit

If  $f(x)$  approaches  $L$  as  $x$  tends toward  $c$  from the left ( $x < c$ ), we write

$$\lim_{x \rightarrow c^-} f(x) = L.$$

Likewise, if  $f(x)$  approaches  $M$  as  $x$  tends toward  $c$  from the right ( $x > c$ ), then

$$\lim_{x \rightarrow c^+} f(x) = M.$$

### Example

Find  $\lim_{x \rightarrow 2^-} f(x)$  and  $\lim_{x \rightarrow 2^+} f(x)$  for the function

$$f(x) = \frac{x^2 + 3}{x - 2}$$

# One-sided Limit

## Example

Find  $\lim_{x \rightarrow -1^-} f(x)$  and  $\lim_{x \rightarrow -1^+} f(x)$  for the function

$$f(x) = \begin{cases} \frac{2}{x-1} & \text{if } x < -1 \\ x^2 - x & \text{if } x \geq -1 \end{cases}$$

# Existence of a Limit

## Theorem

The two-sided limit  $\lim_{x \rightarrow c} f(x)$  exists if and only if the two one-sided limits  $\lim_{x \rightarrow c^-} f(x)$  and  $\lim_{x \rightarrow c^+} f(x)$  both exist and are equal, and then

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c^-} f(x) = \lim_{x \rightarrow c^+} f(x)$$

## Example

Determine whether  $\lim_{x \rightarrow 1} f(x)$  exists, where

$$f(x) = \begin{cases} 2x + 1 & \text{if } x < 1 \\ -x^2 + 2x + 2 & \text{if } x \geq 1 \end{cases}$$

# Continuity

## Continuity

A function  $f$  is *continuous* at  $c$  if all three of these conditions are satisfied:

- $f(c)$  is defined
- $\lim_{x \rightarrow c} f(x)$  exists
- $\lim_{x \rightarrow c} f(x) = f(c)$

If  $f(x)$  is not continuous at  $c$ , it is said to have a *discontinuity* there.

## Example

Decide if  $f(x) = x^3 - x^2 + x - 4$  is continuous at  $x = 0$ .

# Continuity

## Example

Decide if  $f(x) = \frac{2x + 5}{2x - 4}$  is continuous at  $x = 2$ .

# Continuity

## Continuity of Polynomials and Rational Functions

A polynomial or a rational function is continuous *wherever it is defined*.

### Example

List all values of  $x$  for which  $f(x)$  is not continuous

$$f(x) = \frac{x^2 - 2x + 1}{x^2 - x - 2}$$

# Continuity

## Example

Decide if  $f(x) = \begin{cases} x + 1 & \text{if } x < 0 \\ x - 1 & \text{if } x \geq 0 \end{cases}$  is continuous at  $x = 0$ .

# Continuity

## Example

Find the value of the constant  $A$  such that the function

$$f(x) = \begin{cases} 1 - 2x & \text{if } x < 2 \\ Ax^2 + 2x - 3 & \text{if } x \geq 2 \end{cases}$$

will be continuous for all  $x$ .

# Intermediate Value Property

## The intermediate value property

If  $f(x)$  is continuous on the interval  $a \leq x \leq b$  and  $L$  is a number between  $f(a)$  and  $f(b)$ , the  $f(c) = L$  for some number  $c$  between  $a$  and  $b$ . In other words, a continuous function attains all values between any two of its values.

## Example

Show that the equation  $\sqrt[3]{x} = x^2 + 2x - 1$  must have at least one solution on the interval  $0 \leq x \leq 1$ .