



Shorter Notes: The Center of a Convex Set

Teck-Cheong Lim

Proceedings of the American Mathematical Society, Volume 81, Issue 2 (Feb., 1981),
345-346.

Your use of the JSTOR database indicates your acceptance of JSTOR's Terms and Conditions of Use. A copy of JSTOR's Terms and Conditions of Use is available at <http://www.jstor.org/about/terms.html>, by contacting JSTOR at jstor-info@umich.edu, or by calling JSTOR at (888)388-3574, (734)998-9101 or (FAX) (734)998-9113. No part of a JSTOR transmission may be copied, downloaded, stored, further transmitted, transferred, distributed, altered, or otherwise used, in any form or by any means, except: (1) one stored electronic and one paper copy of any article solely for your personal, non-commercial use, or (2) with prior written permission of JSTOR and the publisher of the article or other text.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

Proceedings of the American Mathematical Society is published by American Mathematical Society. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Proceedings of the American Mathematical Society
©1981 American Mathematical Society

JSTOR and the JSTOR logo are trademarks of JSTOR, and are Registered in the U.S. Patent and Trademark Office. For more information on JSTOR contact jstor-info@umich.edu.

©2000 JSTOR

THE CENTER OF A CONVEX SET

TECK-CHEONG LIM

Let X be a Banach space and K a weakly compact convex nonvoid subset with normal structure [1]. Brodskii and Mil'man [1] constructed, using transfinite induction, a "center" of K which is fixed by every isometry mapping K onto K . In this note, we construct a unique "center" for a weakly compact convex nonvoid subset (not necessarily having normal structure) which is fixed by every affine isometry mapping K into K . A similar theorem for weak* compact convex sets is also possible under some additional assumptions.

CONSTRUCTION. Let K be a nonempty weakly compact convex subset of a Banach space. We shall define C_α for all ordinals α by transfinite induction. Set $C_0 = K$. Let β be an ordinal and suppose that C_α has been defined for $\alpha < \beta$ in such a way that (i) each C_α is a nonempty closed convex subset of K and (ii) C_α , $\alpha < \beta$, is decreasing. If β is a limit ordinal, we set $C_\beta = \bigcap_{\alpha < \beta} C_\alpha$. Otherwise, let γ be the predecessor of β and let

$$S_\beta = \left\{ z \in C_\gamma : z = \frac{1}{2}(x + y) \text{ for some } x, y \in C_\gamma \text{ with } \|x - y\| = \frac{1}{2} \text{diam } C_\gamma \right\}.$$

Then we set $C_\beta = \overline{\text{Co}} S_\beta$. Since C_γ is the closed convex hull of its strongly exposed points (see [2]), it is easy to see that if $\text{card } C_\gamma > 1$, C_β contains no strongly exposed points of C_γ and hence is a proper subset of C_γ . If $\text{card } C_\gamma = 1$, $C_\beta = C_\gamma$. It follows that for sufficiently large ordinals δ , C_δ are identical and consist of exactly one point which we call the center of K .

If X is a Banach space such that the dual of every separable subspace of X is separable, and K is a nonempty weak* compact convex subset of X^* , then every weak* closed convex nonempty subset of K is the weak* closed convex hull of its weak* strongly exposed points (see [5]-[8]). With appropriate changes, the prior construction applies to this situation; in particular, replacing C_β by $\overline{\text{Co}}^*(S_\beta)$, where $\overline{\text{Co}}^*$ denotes the weak* closure. Thus K has a unique center.

THEOREM 1. *Let K be a nonempty weakly compact convex subset of a Banach space. The center of K is a fixed point of every affine isometry mapping K into K .*

PROOF. Note that in the construction, each C_α is mapped into itself by every affine isometry of K into K .

Received by the editors April 21, 1980.

1980 *Mathematics Subject Classification.* Primary 46B20; Secondary 47H10.

© 1981 American Mathematical Society
0002-9939/81/0000-0093/\$01.50

THEOREM 2. *Let X be a Banach space such that the dual of every separable subspace of X is separable. Let K be a weak* compact convex nonempty subset of X^* . The center of K is a fixed point of every weak* continuous affine isometry mapping K into K .*

PROOF. If T is an affine isometry, then $T(\text{Co } S_\beta) \subseteq \text{Co } S_\beta$. By the weak* continuity, $T(C_\beta) = T(\overline{\text{Co}^* S_\beta}) \subseteq C_\beta$.

REMARKS. 1. It also follows from the Ryll-Nardzewski fixed point theorem (see [4]) that the family of affine isometries on K has a common fixed point (which is not necessarily the center). Our approach follows that of Namioka-Asplund [4].

2. The assumption of weak* continuity in Theorem 2 cannot be removed since Example 1 in [3] shows that there are fixed point free affine isometries.

REFERENCES

1. M. S. Brodskii and D. P. Mil'man, *On the center of a convex set*, Dokl. Akad. Nauk SSSR **59** (1948), 837–840. MR **9**, 448.
2. J. Diestel and J. J. Uhl, *Vector measures*, Math. Surveys, no. 15, Amer. Math. Soc., Providence, R.I., 1977 (particularly Chapter 7).
3. T. C. Lim, *Asymptotic centers and nonexpansive mappings in some conjugate Banach spaces*, Pacific J. Math. (to appear).
4. I. Namioka and E. Asplund, *A geometric proof of Ryll-Nardzewski's fixed point theorem*, Bull. Amer. Math. Soc. **73** (1967), 443–445. MR **35** #799.
5. I. Namioka and R. R. Phelps, *Banach spaces which are Asplund spaces*, Duke Math. J. **42** (1975), 735–750. MR **52** #11544.
6. C. Stegall, *The Radon-Nikodym property in conjugate Banach spaces*, Trans. Amer. Math. Soc. **206** (1975), 213–223. MR **51** #10581.
7. _____, *The duality between Asplund spaces and spaces with the Radon-Nikodym property*, Israel J. Math. **29** (1978), 408–412.
8. J. J. Uhl, *A note on the Radon-Nikodym theorem for Banach spaces*, Rev. Roumaine Math. Pures Appl. **17** (1972), 113–115.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, ILLINOIS 60637

Current address: Department of Mathematics, George Mason University, Fairfax, Virginia 22030