



On the Normal Structure Coefficient and the Bounded Sequence Coefficient

Teck-Cheong Lim

Proceedings of the American Mathematical Society, Volume 88, Issue 2 (Jun., 1983),
262-264.

Your use of the JSTOR database indicates your acceptance of JSTOR's Terms and Conditions of Use. A copy of JSTOR's Terms and Conditions of Use is available at <http://www.jstor.org/about/terms.html>, by contacting JSTOR at jstor-info@umich.edu, or by calling JSTOR at (888)388-3574, (734)998-9101 or (FAX) (734)998-9113. No part of a JSTOR transmission may be copied, downloaded, stored, further transmitted, transferred, distributed, altered, or otherwise used, in any form or by any means, except: (1) one stored electronic and one paper copy of any article solely for your personal, non-commercial use, or (2) with prior written permission of JSTOR and the publisher of the article or other text.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

Proceedings of the American Mathematical Society is published by American Mathematical Society. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Proceedings of the American Mathematical Society
©1983 American Mathematical Society

JSTOR and the JSTOR logo are trademarks of JSTOR, and are Registered in the U.S. Patent and Trademark Office. For more information on JSTOR contact jstor-info@umich.edu.

©2000 JSTOR

ON THE NORMAL STRUCTURE COEFFICIENT AND THE BOUNDED SEQUENCE COEFFICIENT

TECK-CHEONG LIM

ABSTRACT. The two notions of normal structure coefficient and bounded sequence coefficient introduced by Bynum are shown to be the same. A lower bound for the normal structure coefficient in L^p , $p > 2$, is also given.

Let X be a Banach space and C a closed convex bounded subset of X . For each x in C , let $R(x, C) = \sup\{\|x - y\|: y \text{ in } C\}$ and let $R(C)$ denote the Chebyshev radius of C [2, p. 178]:

$$R(C) = \inf\{R(x, C): x \text{ in } C\}.$$

Let $D(C)$ denote the diameter of C , $D(C) = \sup\{\|x - y\|: x, y \in C\}$.

For a bounded sequence $\{x_n\}$ in X , the asymptotic diameter $A(\{x_n\})$ of $\{x_n\}$ is defined to be $\lim_n \sup\{\|x_k - x_m\|: m \geq n, k \geq n\}$.

In [1], Bynum introduced the following two coefficients of X , called the normal structure coefficient and the bounded sequence coefficient respectively:

$$N(X) = \inf\{D(C)/R(C): C \text{ closed convex bounded nonempty} \\ \text{subsets of } X \text{ with } |C| > 1\},$$

$$BS(X) = \sup\left\{M: \text{for every bounded sequence } \{x_n\} \text{ in } X, \text{ there} \\ \text{exists } y \text{ in } \overline{\text{Co}}(x_n) \text{ such that } M \lim_n \sup\|x_n - y\| \leq A(\{x_n\})\right\}.$$

Another coefficient relating to the asymptotic radius of a sequence (see e.g. [3]) can be defined as follows: Let $\{x_n\}$ be a bounded sequence in X . For each x in X , define

$$r(x, \{x_n\}) = \lim_n \sup\|x_n - x\|.$$

The number $r(\{x_n\}) = \inf\{r(x, \{x_n\}): x \in \overline{\text{Co}}(x_n)\}$ will be called the asymptotic radius of $\{x_n\}$, or more precisely, the asymptotic radius of $\{x_n\}$ w.r.t. $\overline{\text{Co}}(x_n)$. We shall denote the coefficient

$$\inf\{A(\{x_n\})/r(\{x_n\}): \{x_n\} \text{ bounded nonconvergent sequences in } X\}$$

by $A(X)$.

Received by the editors April 13, 1982 and, in revised form, July 23, 1982.

1980 *Mathematics Subject Classification*. Primary 46B20, 47H09, 47H10; Secondary 52A05.

Key words and phrases. Normal structure coefficient, bounded sequence coefficient, Chebyshev radius, asymptotic radius, inequality in L^p .

©1983 American Mathematical Society
 0002-9939/82/0000-0968/\$01.50

In [1], Bynum mentioned that the two coefficients $N(X)$ and $BS(X)$ are equal in a separable Banach space X . In this note, we shall show that the three coefficients are equal in any Banach space X .

THEOREM 1. *For a Banach space X , $N(X) = BS(X) = A(X)$.*

PROOF. It follows readily from the definition that $BS(X) = A(X)$. Indeed, we may assume that the sequences in the definition of $BS(X)$ are nonconvergent. Clearly $BS(X) \leq A(X)$. On the other hand for each $\lambda > 1$, $A(X)/\lambda$ belongs to the defining set of $BS(X)$ and thus $BS(X) \geq A(X)$. Bynum [1] proved that $N(X) \leq BS(X)$. To prove that $BS(X) \leq N(X)$, it suffices to show that for any bounded convex nonempty set C with more than one point, there is a separable closed convex subset C_1 such that $R(C_1) = R(C)$. Indeed, if $\{x_n\}$ is a dense sequence in C_1 and M is a number in the defining set of $BS(X)$, then

$$M \leq A(\{x_n\}) / \limsup_n \|x_n - y\| = \frac{D(C_1)}{R(y, C_1)} \leq \frac{D(C_1)}{R(C_1)} \leq \frac{D(C)}{R(C)}.$$

To construct C_1 , we start out with a sequence of points $\{z_n\}$ in C such that $\lim_{n \rightarrow \infty} R(z_n, C) = R(C)$. Let $U_1 = \text{Co}(\{z_n\})$. Let $V_1 = \{x \in U_1 : R(x, U_1) < R(C)\}$ and let W_1 be a countable dense subset of V_1 . For each x in W_1 , let D_x be a sequence of points in C such that $R(x, D_x) \geq R(C)$. Let X_1 be the countable set $\cup \{D_x : x \in W_1\}$ and $U_2 = \text{Co}(U_1 \cup X_1)$. We define similarly V_2, W_2 and X_2 from U_2 and continue this process to obtain an increasing sequence of convex sets $U_1 \subset U_2 \subset U_3 \subset \dots \subset U_n \subset \dots$. Let $C_1 = \overline{\text{Co}}(\cup U_n)$. C_1 is separable. Since $R(z_n, C_1) \leq R(z_n, C)$ and $\lim_{n \rightarrow \infty} R(z_n, C) = R(C)$, we have $R(C_1) \leq R(C)$. From the way U_n are constructed, $R(x, U_{n+1}) \geq R(C)$ for each $x \in U_n$. It follows that $R(C_1) \geq R(C)$ and the proof is complete.

For $0 < \mu \leq \frac{1}{2}$ and $p > 2$, denote by $x(\mu)$ the unique solution of the equation

$$\lambda x^{p-1} - \mu - (\lambda x - \mu)^{p-1} = 0$$

in the interval $\mu/\lambda \leq x \leq 1$. Define $g(\mu)$, $0 < \mu \leq 1$, by

$$g(\mu) = \lambda \mu \frac{1 + x(\lambda \wedge \mu)^{p-1}}{(1 + x(\lambda \wedge \mu))^{p-1}}$$

where $\lambda = 1 - \mu$. We proved in [4] the following inequality in L^p ($p > 2$):

$$\|\lambda x + \mu y\|^p + g(\mu) \|x - y\|^p \leq \lambda \|x\|^p + \mu \|y\|^p$$

and that

$$\sup_{0 < \mu \leq 1} \frac{g(\mu)}{\mu} = \frac{1 + \alpha^{p-1}}{(1 + \alpha)^{p-1}},$$

where α is the unique solution of

$$(p - 2)x^{p-1} + (p - 1)x^{p-2} - 1 = 0$$

in the interval $0 \leq x \leq 1$.

THEOREM 2. For $X = L^p$, $p > 2$,

$$N(X) \geq \left(1 + \frac{1 + \alpha^{p-1}}{(1 + \alpha)^{p-1}} \right)^{1/p}.$$

PROOF. For a closed convex bounded set C in X , let R and D be the Chebyshev radius and the diameter of C respectively. Let z be the Chebyshev center of C . For x, y in C and $0 < \mu \leq 1$, we have

$$\|\lambda z + \mu y - x\|^p + g(\mu)\|z - y\|^p \leq \lambda\|z - x\|^p + \mu\|y - x\|^p.$$

Taking sup over x in C and noting that $R \leq \sup\{\|\lambda z + \mu y - x\|: x \in C\}$, we obtain

$$R^p + g(\mu)\|z - y\|^p \leq \lambda R^p + \mu \sup\{\|y - x\|^p: x \in C\}.$$

It follows, after taking sup over y in C , that $(\mu + g(\mu))R^p \leq \mu D^p$ and hence

$$\frac{D}{R} \geq \left(1 + \sup_{0 < \mu \leq 1} \frac{g(\mu)}{\mu} \right)^{1/p} = \left(1 + \frac{1 + \alpha^{p-1}}{(1 + \alpha)^{p-1}} \right)^{1/p}.$$

Therefore

$$N(X) \geq \left(1 + \frac{1 + \alpha^{p-1}}{(1 + \alpha)^{p-1}} \right)^{1/p}. \quad \square$$

REMARK 1. For $p = 3$ and 4 , we have $\alpha = \sqrt{2} - 1$ and $1/2$ and hence

$$\left(1 + \frac{1 + \alpha^{p-1}}{(1 + \alpha)^{p-1}} \right)^{1/p} = (3 - \sqrt{2})^{1/3} \text{ and } (4/3)^{1/4}$$

respectively.

REFERENCES

1. W. L. Bynum, *Normal structure coefficients for Banach spaces*, Pacific J. Math. **86** (1980), 427–436.
2. R. B. Holmes, *A course in optimization and best approximation*, Lecture Notes in Math., vol. 257, Springer-Verlag, Berlin and New York, 1972.
3. T. C. Lim, *Characterizations of normal structure*, Proc. Amer. Math. Soc. **43** (1974), 313–319.
4. _____, *Fixed point theorems for uniformly Lipschitzian mappings in L^p spaces*, J. Nonlinear Anal. Theory, Method and Appl. (to appear).

DEPARTMENT OF MATHEMATICS, GEORGE MASON UNIVERSITY, 4400 UNIVERSITY DRIVE, FAIRFAX, VIRGINIA 22030